

spenito is required for sex determination in *Drosophila melanogaster*

Dong Yan^a and Norbert Perrimon^{a,b,1}

^aDepartment of Genetics, Harvard Medical School, Boston, MA 02115; and ^bHoward Hughes Medical Institute, Harvard Medical School, Boston, MA 02115

Contributed by Norbert Perrimon, August 12, 2015 (sent for review May 4, 2015; reviewed by Brian Oliver)

***Sex-lethal (Sxl)* encodes the master regulator of the sex determination pathway in *Drosophila* and acts by controlling sex identity in both soma and germ line. In females *Sxl* maintains its own expression by controlling the alternative splicing of its own mRNA. Here, we identify a novel sex determination gene, *spenito (nito)* that encodes a SPEN family protein. Loss of *nito* activity results in stem cell tumors in the female germ line as well as female-to-male somatic transformations. We show that *Nito* is a ubiquitous nuclear protein that controls the alternative splicing of the *Sxl* mRNA by interacting with *Sxl* protein and pre-mRNA, suggesting that it is directly involved in *Sxl* auto-regulation. Given that SPEN family proteins are frequently mutated in cancers, our results suggest that these factors might be implicated in tumorigenesis through splicing regulation.**

germ-line stem cell | sex determination | alternative splicing

Sex determination in *Drosophila* is under the control of the master regulatory gene *Sex-lethal (Sxl)* (1). *Sxl* acts downstream of the X-chromosome counting mechanism and encodes a female-specific RNA binding protein. Once activated, *Sxl* maintains its own expression by regulating the alternative splicing of its pre-mRNA. *Sxl* controls female fate by controlling somatic and germ-line sex identity as well as dosage compensation (2). In female somatic cells, *Sxl* controls the alternative splicing of *transformer (tra)*, which together with *transformer2 (tra2)* controls the alternative splicing of *doublesex (dsx)* and *fruitless (fru)*. *dsx* and *fru* in turn encode sex-specific transcription factors that control male versus female morphology, physiology, and behavior (3, 4). In addition, *Sxl* represses the male-specific dosage compensation system by regulating *male-specific lethal 2 (msl-2)* both at the level of alternative splicing and translational control (5).

In the female germ-line *Sxl* regulates sex identity by a different mechanism, as *tra*, *tra2*, *msl-2* have no roles in the germ line (2). In the ovary, germ-line stem cells (GSCs) located at the anterior tip of the germarium divide to produce another GSC and a cystoblast (CB) that is committed to differentiate. *Sxl* protein accumulates to high levels in the GSCs/CBs and is required for the proper differentiation of the germ cells (6). Germ cells lacking *Sxl* cannot differentiate and instead produce stem cell tumors. The identity of *Sxl* target genes in the germ line is not well characterized; however, a recent study indicates that *nanos*, a gene required for GSC maintenance, is a *Sxl* target (7). Indeed, *Sxl* has been proposed to promote the differentiation of GSCs by downregulating *Nanos* levels in CBs by binding to the *nanos* 3' UTR (7). In addition, *Sxl* is also important for repressing the expression of testis-specific genes, including *Phf7*, a male germ-line identity gene (8). In the absence of *Sxl*, *Phf7* is mis-expressed leading to germ-line tumors (9).

Sxl does not act alone to control splicing. Several genes, including *sans fille (snf)*, *virilizer (vir)*, *female-lethal-2-d (fl(2)d)*, *SPF45*, *U1-70K*, *U2af38*, *U2af50*, and *protein partner of sans-fille (pps)* facilitate *Sxl* splicing autoregulation (10–18). Except for *pps*, these genes encode either general splicing factors or proteins associated with spliceosomes. They all act to maintain the *Sxl* autoregulatory splicing loop by interacting with *Sxl* itself. In addition, some of them are involved in the splicing of other *Sxl* splicing targets such as *tra* or *msl-2* (1). Interestingly, these genes

have essential functions besides *Sxl* regulation and null mutations are associated with zygotic lethality in both sexes. Therefore, the roles of these factors in sex determination were revealed from genetic interactions (*snf*, *U1-70K*, *U2af38*) (11, 17), temperature-sensitive mutation (*vir*) (15), clonal analysis (*fl(2)d)* (10), or biochemical studies (*SPF45*, *U2af50* and *pps*) (12, 14, 17).

Here, we characterize *spenito (nito)*, a novel regulator of *Sxl*, which is required to maintain sex identity and *Sxl* levels in both the female germ-line and somatic tissues. *Nito* is required for the proper alternative splicing of the *Sxl* pre-mRNA in both germ line and soma, and forms a complex with *Sxl* protein and its pre-mRNA, thus identifying an important component of the sex determination pathway.

Results

nito Is an Essential Gene Required for Ovarian GSC Differentiation.

nito was identified from our previous RNAi screen in *Drosophila* GSCs (19). Specifically, RNAi knockdown of *nito* driven by the germ-line-specific *MTD-Gal4* driver resulted in complete sterility in females. In wild-type ovarioles, two or three GSCs are located in the anterior tip of the germarium (Fig. 1 *A–B'*). Strikingly, *nito* shRNA ovarioles are filled with undifferentiated stem-cell-like cells, and nurse cells and oocytes are not formed (Fig. 1 *C* and *C'*, compared with WT in Fig. 1 *B* and *B'*). Further, stem-cell-like cells associated with *nito* shRNA ovaries retain their proliferative potential as shown by staining with the mitotic marker phosphorylated histone H3 (pH3) (Fig. 1*E*, compared with WT in Fig. 1*D*). Note that the same stem-cell tumor phenotype was observed with three independent *nito* shRNAs and two long dsRNA RNAi lines (*Methods*) (Fig. S1 *A–B'*), indicating that *nito* is an essential gene required for GSC differentiation.

***nito* Is Required for Sex Determination in the Soma.** Because the germ-line phenotype of *nito* could reflect perturbations in a number of developmental processes affecting either germ-line

Significance

Sex determination is a fundamental biological problem faced by all metazoans. To understand the sex determination pathway, it is important to identify all the genes involved in this process. In this study, we have identified a novel gene, *spenito (nito)*, which is required for sex determination in *Drosophila melanogaster*. Loss of *nito* function in the soma transforms female tissues to male, and loss of *nito* function in female germ-line stem cells changes their sexual identity and prevents them from proper differentiation. We show that *nito* is a cofactor for *Sex-lethal (Sxl)* auto regulation, a process that remains an important textbook model for regulated alternative splicing.

Author contributions: D.Y. and N.P. designed research; D.Y. performed research; D.Y. and N.P. analyzed data; and D.Y. and N.P. wrote the paper.

Reviewers included: B.O., NIH.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. Email: perrimon@receptor.med.harvard.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1515891112/-DCSupplemental.




Fig. 2. Nito is required for sex determination in somatic tissues. (A and B) *dome-Gal4* drives expression in the first pair of leg discs (A) and genital discs (B), as shown by *UAS-2xEYFP*. (C) Foreleg of a WT male with the dark thickened sex comb bristles. (D) Foreleg of a WT female. Note the absence of sex combs. (E) Foreleg of a female fly expressing *nito* shRNA driven by *dome-Gal4*. Some bristles are transformed into male sex combs. (F and G) Genitalia of wild-type male (F) and female (G) flies showing distinct morphology, such as penis apparatus and claspers in male (F, arrow and arrowhead, respectively) and vaginal bristles in female (G, arrow). (H) *nito* shRNA driven by *dome-Gal4* transforms female genital morphology into male-like, as evidenced by the absence of vaginal bristles and appearance of structures resembling penis apparatus (arrow) and claspers (arrowhead).

a deficiency of the *nito* locus indicating that the lethality is likely due to the *nito* mutation. Consistent with this, *nito* RNAi driven by a ubiquitous Gal4 such as *actin-Gal4* or *tubulin-Gal4* is associated with larval lethality.

Nito Regulates Sxl Levels by Controlling Sxl Alternative Splicing. The phenotype associated with loss of Nito function in both soma and germ line suggests that Nito may regulate *Sxl* activity. In ovaries, *Sxl* is enriched in GSCs and their immediate daughter cells (Fig. 4E) (6), whereas in somatic tissues such as wing discs *Sxl* is expressed ubiquitously in females but absent in males (Fig. 4 A and B). We expressed *nito* shRNA in the germ line using *MTD-Gal4* and in the dorsal half of the wing disk using *ap-Gal4*. Strikingly, knockdown of *nito* in both tissues led to a significant reduction of *Sxl* levels (Fig. 4 F, C, and C'). Note that similar results were obtained using two additional *nito* shRNA lines as well as in homozygous *nito*¹ mutant clones (Fig. 4 D and D' and Fig. S2 A' and B'). However, the level of *Sxl* is not affected when Split-ends (Spen), another member of the SPEN family, was knocked down by shRNA in the wing disk (Fig. S2C), indicating a specific role of Nito in the sex determination pathway. Altogether, we have identified a new component of the *Drosophila*

sex determination pathway that acts in both the germ line and soma by affecting *Sxl* levels.

Sxl transcripts are alternatively spliced, with exon 3 containing a stop codon that is included in males but skipped in females, leading to truncated *Sxl* forms in males but functional proteins in females (Fig. 4G) (22). Because Nito has three RRM domains, we asked whether Nito has a potential role in *Sxl* splicing and used a pair of primers that detects the small female and large male spliced *Sxl* products (Fig. 4G) (12) to analyze *Sxl* splicing in the absence of Nito. In *nito* shRNA female wing discs or ovaries, a large band corresponding to the male-specific spliced form was clearly detected (Fig. 4G). Note that in wing discs and ovaries, female-specific transcripts are detected due to contributions from remaining WT disk cells or somatic follicle cells, respectively, as *nito* was knocked down in half of the discs or in the germ line only. Together, our results indicate that Nito regulates *Sxl* levels by controlling its alternative splicing.

Nito Interacts with Sxl Protein and its Pre-mRNA in S2 Cells. We then analyzed how Nito controls *Sxl* alternative splicing. Because the key protein that binds *Sxl* pre-mRNA and inhibits splicing of male-specific exon 3 is *Sxl* itself (2), we examined whether Nito interacts with *Sxl* using a coimmunoprecipitation (co-IP) assay in *Drosophila* S2 cells. GFP-Nito and HA-*Sxl* were expressed either individually or in combination in S2 cells, and GFP alone was used as a control (Fig. 5A). HA-*Sxl* was detected in the precipitate obtained using anti-GFP nanobody from GFP-Nito cells, but not from GFP-expressing cells. Similarly, GFP-Nito, but not GFP, was pulled down by HA-*Sxl*. Further, we tested whether the interaction between Nito and *Sxl* is dependent on the presence of RNA. Interestingly, the amount of *Sxl*-HA pulled down by GFP-Nito is strongly reduced in the presence of RNase, suggesting that the Nito/*Sxl* interaction is mediated or stabilized by RNA (Fig. 5B). Finally, we performed RNA immunoprecipitation (RIP) experiments in S2 cells to analyze whether Nito can interact with *Sxl* pre-mRNA, which was detected by RT-PCR using an intron 3-exon 4 primer pair (12). As shown in Fig. 5C, GFP-*Sxl*, GFP-Nito, but not GFP alone, can pull down *Sxl* pre-mRNA from cell lysates. Together, the specific interactions between Nito, *Sxl*, and *Sxl* pre-mRNA support the model that Nito forms a complex with *Sxl* and that they together regulate alternative splicing of *Sxl* mRNA (Fig. 5D).

Screen for Additional Splicing Genes Involved in Sex Determination.

Because we identified *nito* as an important gene involved in sex determination by regulating *Sxl* splicing, we asked whether there are other unidentified genes in the *Drosophila* genome acting in this pathway. Because most known *Sxl*-autoregulatory proteins are associated with spliceosomes (2), we screened a collection of 316 RNAi lines representing 247 splicing-associated genes and RNA-binding proteins (Dataset S1), using *MTD-Gal4* for the germ line and *dome-Gal4* for somatic phenotypes. In addition, we also included in our screen a wing-specific driver, *nub-Gal4*, as many RNAi lines exhibit lethality with *dome-Gal4* and prevent the detection of potential sex determination phenotypes. Because *Sxl* is responsible for the larger wing size in females, examination of wing size using *nub-Gal4* allows detection of potential genes involved in sex determination, as shown in the case of *nito* (Fig. S3 A-F).

Our screen successfully identified known components of the sex determination pathway: *tra* and *tra2* RNAi showed strong female-to-male transformation in sex combs and genitalia when induced by *dome-Gal4*; and *jl(2)d* and *vir* were identified using the *nub-Gal4* driver as both have a stronger effect in female wings than in male wings. Strikingly, besides these genes we did not identify any other genes that showed a sex-related phenotype. However, we did characterize GSC differentiation phenotypes associated with *tsu*, *mago*, *RnpS1*, *Rbp9* RNAi lines, as well




Fig. 3. Nito is a ubiquitous nuclear protein that shows no sex-biased expression. (A) Schematic diagram showing domain structures of Spen and Nito proteins and the peptide used to generate the Nito antibody. (B) *nito* mRNA levels in male and female WT wing discs were measured by qRT-PCR. Error bars represent SDs. (C) Nito antibody recognizes a single band of the predicted ~89 kDa size in S2 cell lysates. (D–F) Nito antibody stainings in WT female (D) or male (E) wing discs and ovarioles (F). (G–G') Expression of *Sxl* shRNA in the dorsal half of the wing disk (below the dashed line) using *ap-Gal4* leads to depletion of *Sxl* protein, but has no effect on Nito protein levels. (H–H') Nito antibody staining in wing discs containing *nito*¹ mutant clones, marked by the absence of GFP. Note the absence of Nito staining in *nito*¹ clones.

as wing growth and pattern defects with *kul*, *CG7879*, *ASPP*, *tst* and *Syp* RNAi lines. These data provide a valuable resource of phenotypes associated with splicing-related genes (data available at www.flyrnai.org/RSPV.html).

Discussion

We describe the characterization of Nito as a novel component of the *Drosophila* sex determination pathway. *Nito* loss-of-function results in stem-cell tumor phenotypes in the germ-line and sexual transformations in the soma. Interestingly, Nito affects *Sxl* protein levels in both GSCs and somatic tissues by regulating *Sxl* pre-mRNA alternative splicing, most likely directly as Nito interacts with the *Sxl* protein and pre-mRNA. The role of Nito is reminiscent of the previously reported roles of splicing factors in *Sxl* auto regulation, such as both subunits of U2AF (17), U1-70K

(17, 18), Fl(2)d (10, 23), SPF45 (13, 14), Vir (15, 24), and Snf (11, 16). Our data support earlier reports that *Sxl* physically interacts with components of the spliceosome to simultaneously block utilization of the 3' and 5' splice sites of the male exon.

Nito and Spen are members of the SPEN protein family that are evolutionarily conserved from plants, worms, flies to mice and humans (25). Both proteins contain three N-terminal RRM domains and one C-terminal SPOC domain. The sequence similarity between these domains is low and there is no conservation outside these motifs, suggesting that they have evolved specific functions following a duplication event (20), as indicated by our observation that *spen* is not required for *Sxl* regulation (Fig. S2C). In *Drosophila*, *spen* was first identified in several genetic screens looking for components of the receptor tyrosine kinase (RTK) signaling pathway (26). Subsequent studies found




Fig. 4. Nito is required for Sxl levels and regulates Sxl mRNA splicing. (A and B) Sxl stainings in WT male (A) and female (B) wing discs. (C–C') Expressing *nito* shRNA in the dorsal half of the disk (below the dashed line) using *ap-Gal4* leads to strong reduction of Nito (C) and Sxl (C') stainings. (D–D') Sxl antibody staining (D') in wing discs containing *nito*¹ mutant clones, which are marked by the absence of GFP (D). Note the absence of Nito and Sxl staining in *nito*¹ clones. (E and F) Sxl stainings in WT egg chambers (E) or in egg chambers expressing *nito* shRNA by *MTD-Gal4* (F). Scale bars: 20 μ m. (G) Diagram showing the alternative splicing event that produces the male- or female-specific Sxl transcripts. The arrows indicate the primers used for RT-PCR. Sxl splicing was analyzed by RT-PCR using RNA extracted from wing discs or ovaries. Male-specific bands: 2–3–4. Female-specific bands: 2–4.

that *spen* is implicated in a variety of cellular and developmental processes including neuronal cell fate specification, axon guidance, cell cycle, Hox gene regulation, and cell death (27–30). These pleiotropic effects are likely due to the involvement of *spen* in multiple signaling pathways (31–33). However, the molecular mechanisms underlying the function of Spen in these pathways are not understood.

Genetic studies in *Drosophila* have shown that *nito* over-expression results in a rough eye phenotype (20) and that it plays a redundant role with *spen* in Wnt signaling (21), but how Nito is involved in these processes is not known. Biochemical studies indicate that Nito, like its human ortholog, copurify with the precatalytic spliceosome (complex B) (34). In addition, *nito*, as well as many other splicing factors, was identified in an RNAi screen for RAS/MAPK signaling components (35). Consistent with these findings, we find that *nito* is required for the alternative splicing of the master sex-determination gene *Sxl*. Previously, both Spen and Nito were thought to act mainly as transcription factors through their SPOC domains, our findings however clearly indicate that Nito is involved in mRNA splicing. It is intriguing to note that PPS, another important factor required for *Sxl* splicing, also has a SPOC domain (12). Similar to Nito, PPS also forms a complex with Sxl protein and its pre-mRNA (12). In the future it will be crucial to dissect how different protein domains contribute to the function of SPEN family proteins.

Then what is the “main” role of *nito*? On one hand, the phenotypes in the sex comb, genitalia and germ line appear specific to *Sxl* and such phenotypes do not depend on the genetic interaction with other genes in the sex determination pathway. On the other hand, *nito* clearly has other non-sex-specific functions, as revealed by the lethality, rough eye, and wing phenotype observed in both sexes (Fig. S3 A–F). Because a null allele of *nito* is associated with zygotic lethality, the RNAi knockdown approach is a powerful method to reveal sex-related phenotypes. Interestingly, our RNAi screen targeting splicing factors did not identify any new additional sex determination genes, indicating that there are a limited number of genes yet to be identified in this pathway. Finally, intriguingly, three recent studies have identified SPEN and Rbm15 (the mouse and human ortholog of Nito) as factors interacting with *Xist*, the long noncoding RNA that is essential for dosage compensation in mammals (36–38). Clearly, future experiments such as RNA-seq will be necessary to elucidate the mechanism and logic of Nito-mediated signaling events.

Rbm15, also known as OTT, was originally identified from infants with acute megakaryoblastic leukemia (AMKL) (39, 40). The t(1, 22) chromosomal translocation results in fusion of




Fig. 5. Nito interacts with Sxl and Sxl pre-mRNA in S2 cells. (A) HA-Sxl, GFP-Nito or GFP expression vectors were transfected individually or together into *Drosophila* S2 cells. Cell lysates were immunoprecipitated using GFP nanobody or anti-HA antibody and analyzed by Western blot. GFP alone is used as a control. Asterisk indicates IgG heavy chain. (B) S2 cells were transfected with GFP-Nito and HA-Sxl, and Co-IP was performed using GFP nanobody in the absence or presence of RNase A and RNase T1. (C) S2 cells were transfected with GFP-Nito, GFP-Sxl or GFP and immunoprecipitated with GFP nanobody. The presence of Sxl pre-mRNA was detected by RT-PCR using an intron 3/exon 4 primer pair. GFP-Sxl was used as a positive control and GFP alone as a negative control. (D) Model: Nito forms a complex with Sxl and together they repress the splicing of Sxl exon 3 in female tissues.

RBM15 and MKL1, and the fusion protein is responsible for AMKL development as shown in a mouse model (41). In addition to this chromosome translocation, recent cancer genome sequencing projects have found that RBM15 and SPEN (also known as SHARP) are mutated in many different types of cancers, such as adenoid cystic carcinomas and bladder cancers (42). Given that SPEN family proteins are frequently mutated or deleted in cancers, they have been proposed to act as potential tumor suppressors (42). Studies of Spen and Nito in *Drosophila* will provide mechanistic insights to our understanding of this important family of proteins.

Methods

Details on the fly strains used in this study, as well as how the null *nito* mutation was isolated and how *nito* clones were generated can be found in *SI Methods*. Protocols used for antibody staining, reagents, how Nito antibodies were generated, coimmunoprecipitation protocols, RT-PCR, and information on primers and RNA immunoprecipitation (RIP), can be found in *SI Methods*.

ACKNOWLEDGMENTS. We thank Ken Cadigan, Jamila Horabin, Ilaria Rebay, and Helen Salz for fly stocks and antibodies. We thank Mitzi Kuroda, Leonard Rabinow, Stephanie Mohr, Richard Binari, Benjamin Housden, and Ben Ewen-Campen for comments on the manuscript. D.Y. is supported by Damon Runyon Cancer Research and Charles A. King Trust Postdoctoral Fellowship. N.P. is an investigator of the Howard Hughes Medical Institute.

- Cline TW, Meyer BJ (1996) Vive la différence: Males vs females in flies vs worms. *Annu Rev Genet* 30:637–702.
- Salz HK, Erickson JW (2010) Sex determination in *Drosophila*: The view from the top. *Fly (Austin)* 4(1):60–70.
- Clough E, et al. (2014) Sex- and tissue-specific functions of *Drosophila* doublesex transcription factor target genes. *Dev Cell* 31(6):761–773.
- Ryner LC, et al. (1996) Control of male sexual behavior and sexual orientation in *Drosophila* by the fruitless gene. *Cell* 87(6):1079–1089.
- Camara N, Whitworth C, Van Doren M (2008) The creation of sexual dimorphism in the *Drosophila* soma. *Curr Top Dev Biol* 83:65–107.
- Chau J, Kulnane LS, Salz HK (2009) Sex-lethal facilitates the transition from germline stem cell to committed daughter cell in the *Drosophila* ovary. *Genetics* 182(1):121–132.
- Chau J, Kulnane LS, Salz HK (2012) Sex-lethal enables germline stem cell differentiation by down-regulating Nanos protein levels during *Drosophila* oogenesis. *Proc Natl Acad Sci USA* 109(24):9465–9470.
- Yang SY, Baxter EM, Van Doren M (2012) Phf7 controls male sex determination in the *Drosophila* germline. *Dev Cell* 22(5):1041–1051.
- Shapiro-Kulnane L, Smolko AE, Salz HK (2015) Maintenance of *Drosophila* germline stem cell sexual identity in oogenesis and tumorigenesis. *Development* 142(6):1073–1082.
- Granadino B, Campuzano S, Sánchez L (1990) The *Drosophila melanogaster* *fl(2)d* gene is needed for the female-specific splicing of Sex-lethal RNA. *EMBO J* 9(8):2597–2602.
- Oliver B, Perrimon N, Mahowald AP (1988) Genetic evidence that the *sans fille* locus is involved in *Drosophila* sex determination. *Genetics* 120(1):159–171.
- Johnson ML, Nagengast AA, Salz HK (2010) PPS, a large multidomain protein, functions with sex-lethal to regulate alternative splicing in *Drosophila*. *PLoS Genet* 6(3):e1000872.
- Chaouki AS, Salz HK (2006) *Drosophila* SPF45: A bifunctional protein with roles in both splicing and DNA repair. *PLoS Genet* 2(12):e178.
- Lallena MJ, Chalmers KJ, Llamazares S, Lamond AI, Valcárcel J (2002) Splicing regulation at the second catalytic step by Sex-lethal involves 3' splice site recognition by SPF45. *Cell* 109(3):285–296.
- Hilfiker A, Amrein H, Dübendorfer A, Schneider R, Nöthiger R (1995) The gene *virilizer* is required for female-specific splicing controlled by Sxl, the master gene for sexual development in *Drosophila*. *Development* 121(12):4017–4026.
- Flickinger TW, Salz HK (1994) The *Drosophila* sex determination gene *snf* encodes a nuclear protein with sequence and functional similarity to the mammalian U1A snRNP protein. *Genes Dev* 8(8):914–925.
- Nagengast AA, Stitzinger SM, Tseng CH, Mount SM, Salz HK (2003) Sex-lethal splicing autoregulation in vivo: Interactions between SEX-LETHAL, the U1 snRNP and U2AF underlie male exon skipping. *Development* 130(3):463–471.
- Salz HK, et al. (2004) The *Drosophila* U1-70K protein is required for viability, but its arginine-rich domain is dispensable. *Genetics* 168(4):2059–2065.
- Yan D, et al. (2014) A regulatory network of *Drosophila* germline stem cell self-renewal. *Dev Cell* 28(4):459–473.
- Jemc J, Rebay I (2006) Characterization of the split ends-like gene *spenito* reveals functional antagonism between SPOC family members during *Drosophila* eye development. *Genetics* 173(1):279–286.
- Chang JL, Lin HV, Blauwkamp TA, Cadigan KM (2008) *Spenito* and *Split ends* act redundantly to promote *Wingless* signaling. *Dev Biol* 314(1):100–111.
- Bell LR, Horabin JI, Schedl P, Cline TW (1991) Positive autoregulation of sex-lethal by alternative splicing maintains the female determined state in *Drosophila*. *Cell* 65(2):229–239.
- Penn JK, et al. (2008) Functioning of the *Drosophila* Wilms'-tumor-1-associated protein homolog, *Fl(2)d*, in Sex-lethal-dependent alternative splicing. *Genetics* 178(2):737–748.
- Schultz C, Hilfiker A, Nöthiger R (1998) *virilizer* regulates Sex-lethal in the germline of *Drosophila melanogaster*. *Development* 125(8):1501–1507.
- Sánchez-Pulido L, Rojas AM, van Wely KH, Martínez-A C, Valencia A (2004) SPOC: A widely distributed domain associated with cancer, apoptosis and transcription. *BMC Bioinformatics* 5:91.
- Rebay I, et al. (2000) A genetic screen for novel components of the Ras/Mitogen-activated protein kinase signaling pathway that interact with the *yan* gene of *Drosophila* identifies split ends, a new RNA recognition motif-containing protein. *Genetics* 154(2):695–712.
- Kuang B, Wu SC, Shin Y, Luo L, Kolodziej P (2000) split ends encodes large nuclear proteins that regulate neuronal cell fate and axon extension in the *Drosophila* embryo. *Development* 127(7):1517–1529.
- Chen F, Rebay I (2000) split ends, a new component of the *Drosophila* EGF receptor pathway, regulates development of midline glial cells. *Curr Biol* 10(15):943–946.
- Wiellette EL, et al. (1999) *spen* encodes an RNP motif protein that interacts with Hox pathways to repress the development of head-like sclerites in the *Drosophila* trunk. *Development* 126(23):5373–5385.
- Mace K, Tugores A (2004) The product of the split ends gene is required for the maintenance of positional information during *Drosophila* development. *BMC Dev Biol* 4:15.
- Doroquez DB, Orr-Weaver TL, Rebay I (2007) Split ends antagonizes the Notch and potentiates the EGFR signaling pathways during *Drosophila* eye development. *Mech Dev* 124(9-10):792–806.
- Lin HV, et al. (2003) Splits ends is a tissue/promoter specific regulator of *Wingless* signaling. *Development* 130(14):3125–3135.
- Querenet M, Goubard V, Chatelain G, Davoust N, Mollereau B (2015) Spen is required for pigment cell survival during pupal development in *Drosophila*. *Dev Biol* 402(2):208–215.
- Herold N, et al. (2009) Conservation of the protein composition and electron microscopy structure of *Drosophila melanogaster* and human spliceosomal complexes. *Mol Cell Biol* 29(1):281–301.
- Ashton-Beaucage D, et al. (2014) A functional screen reveals an extensive layer of transcriptional and splicing control underlying RAS/MAPK signaling in *Drosophila*. *PLoS Biol* 12(3):e1001809.
- Moindrot B, et al. (2015) A Pooled shRNA screen identifies Rbm15, Spen, and Wtap as factors required for Xist RNA-mediated silencing. *Cell Reports* 12(4):562–572.
- Monfort A, et al. (2015) Identification of Spen as a crucial factor for Xist function through forward genetic screening in haploid embryonic stem cells. *Cell Reports* 12(4):554–561.
- McHugh CA, et al. (2015) The Xist lncRNA interacts directly with SHARP to silence transcription through HDAC3. *Nature* 521(7551):232–236.
- Ma Z, et al. (2001) Fusion of two novel genes, RBM15 and MKL1, in the t(1;22)(p13;q13) of acute megakaryoblastic leukemia. *Nat Genet* 28(3):220–221.
- Mercher T, et al. (2001) Involvement of a human gene related to the *Drosophila* *spen* gene in the recurrent t(1;22) translocation of acute megakaryocytic leukemia. *Proc Natl Acad Sci USA* 98(10):5776–5779.
- Mercher T, et al. (2009) The OTT-MAL fusion oncogene activates RBPJ-mediated transcription and induces acute megakaryoblastic leukemia in a knockin mouse model. *J Clin Invest* 119(4):852–864.
- Su H, et al. (2015) Split end family RNA binding proteins: Novel tumor suppressors coupling transcriptional regulation with RNA processing. *Cancer Translational Medicine* 1(1):21–25.
- Owusu-Ansah E, Banerjee U (2009) Reactive oxygen species prime *Drosophila* haematopoietic progenitors for differentiation. *Nature* 461(7263):537–541.
- Horabin JI (2005) Splitting the Hedgehog signal: sex and patterning in *Drosophila*. *Development* 132(21):4801–4810.
- Stitzinger SM, Conrad TR, Zachlin AM, Salz HK (1999) Functional analysis of SNF, the *Drosophila* U1A/U2B" homolog: identification of dispensable and indispensable motifs for both snRNP assembly and function in vivo. *RNA* 5(11):1440–1450.
- Zhang F, et al. (2012) UAP56 couples piRNA clusters to the perinuclear transposon silencing machinery. *Cell* 151(4):871–884.
- Staller MV, et al. (2013) Depleting gene activities in early *Drosophila* embryos with the "maternal-Gal4-shRNA" system. *Genetics* 193(1):51–61.