

# Control of Proinflammatory Gene Programs by Regulated Trimethylation and Demethylation of Histone H4K20

Joshua D. Stender,<sup>1</sup> Gabriel Pascual,<sup>1</sup> Wen Liu,<sup>2</sup> Minna U. Kaikkonen,<sup>1,3</sup> Kevin Do,<sup>1</sup> Nathanael J. Spann,<sup>1</sup> Michael Boutros,<sup>4</sup> Norbert Perrimon,<sup>4,5</sup> Michael G. Rosenfeld,<sup>2,5</sup> and Christopher K. Glass<sup>1,2,\*</sup>

<sup>1</sup>Department of Cellular and Molecular Medicine

<sup>2</sup>Department of Medicine

University of California, San Diego, La Jolla, CA 92093, USA

<sup>3</sup>A.I. Virtanen Institute, Department of Biotechnology and Molecular Medicine, University of Eastern Finland, P.O. Box 1627, FIN-70120 Kuopio, Finland

<sup>4</sup>Department of Genetics, Harvard Medical School, Boston, MA 02115, USA

<sup>5</sup>Howard Hughes Medical Institute

\*Correspondence: [ckg@ucsd.edu](mailto:ckg@ucsd.edu)

<http://dx.doi.org/10.1016/j.molcel.2012.07.020>

## SUMMARY

Regulation of genes that initiate and amplify inflammatory programs of gene expression is achieved by signal-dependent exchange of coregulator complexes that function to read, write, and erase specific histone modifications linked to transcriptional activation or repression. Here, we provide evidence for the role of trimethylated histone H4 lysine 20 (H4K20me<sub>3</sub>) as a repression checkpoint that restricts expression of toll-like receptor 4 (TLR4) target genes in macrophages. H4K20me<sub>3</sub> is deposited at the promoters of a subset of these genes by the SMYD5 histone methyltransferase through its association with NCoR corepressor complexes. Signal-dependent erasure of H4K20me<sub>3</sub> is required for effective gene activation and is achieved by NF- $\kappa$ B-dependent delivery of the histone demethylase PHF2. Liver X receptors antagonize TLR4-dependent gene activation by maintaining NCoR/SMYD5-mediated repression. These findings reveal a histone H4K20 trimethylation/demethylation strategy that integrates positive and negative signaling inputs that control immunity and homeostasis.

## INTRODUCTION

The survival of vertebrate organisms relies on innate and adaptive immune mechanisms to detect, combat, and eliminate foreign pathogens. Recognition of foreign pathogens by pattern recognition receptors (PRRs) triggers transcriptional activation of genes that amplify inflammatory responses, enable antimicrobial activities, and initiate the development of acquired immunity (Kawai and Akira, 2010; Takeuchi and Akira, 2010). The conservation of key signaling pathways required for these responses,

including the Toll, JAK/STAT, and Immune Deficiency (Imd) pathways, throughout metazoan evolution underscores their importance in effectively defending against bacterial and viral infection (Lemaitre and Hoffmann, 2007; Martinelli and Reichhart, 2005). Macrophages represent a crucial cell type for initial recognition of pathogens based on their expression of numerous PRRs, including the toll-like receptors (TLRs). The TLR family is comprised of a highly conserved set of membrane receptors that recognize specific components of bacterial and viral pathogens including lipoproteins (TLR1/2/6), lipopolysaccharide (TLR4), flagellin (TLR5), single-stranded RNA (TLR7/8), double-stranded RNA (TLR3), and double stranded DNA (TLR9) (Kawai and Akira, 2010; Martinelli and Reichhart, 2005). Upon TLR ligation by their respective ligands, adaptor proteins are recruited to the cytoplasmic domains of TLRs and initiate signaling cascades, which ultimately result in the activation of signal-dependent transcription factors such as nuclear factor- $\kappa$  B (NF- $\kappa$ B), activator protein 1 (AP1), and interferon regulatory factor (IRFs). These factors function in a combinatorial manner to promote the induction of inflammatory cytokines, type I interferons, and numerous other regulatory and effector molecules required for host defense (Kawai and Akira, 2010).

Although required for effective immunity, persistent or inappropriate activation of inflammatory gene programs can contribute to chronic diseases that include type II diabetes (Hotamisligil, 2006), atherosclerosis (Tedgui and Mallat, 2006), cancer (Karin et al., 2006), and neurodegenerative diseases (Glass et al., 2010). Precise control of innate and adaptive immune responses is therefore critical for maintaining proper tissue homeostasis. Multiple mechanisms have been identified that function to prevent signal-independent activation of inflammatory responses and to mediate their resolution upon eradication of the inciting stimulus. Several studies have suggested that many highly inducible inflammatory response genes are maintained in a repressed state under basal conditions by corepressor complexes containing nuclear receptor corepressor 1 (NCoR), silencing mediator of retinoic acid and thyroid hormone receptor (SMRT) complexes, and/or the corepressor of REST

(CoREST) (Ghisletti et al., 2009; Hargreaves et al., 2009; Hoberg et al., 2006; Huang et al., 2009; 2011; Ogawa et al., 2004; Pascual et al., 2005; Saijo et al., 2009; Venteclef et al., 2010). These complexes are actively removed in response to proinflammatory signals as a prerequisite to transcriptional activation and thereby serve as transcriptional checkpoints that regulate the transition from basal to activated states (Ghisletti et al., 2009; Huang et al., 2009, 2011; Ogawa et al., 2004; Pascual et al., 2005; Saijo et al., 2009). Several members of the nuclear receptor family of transcription factors, including glucocorticoid receptors, peroxisome proliferator-activated receptors (PPARs), and liver X receptors (LXRs), exert potent anti-inflammatory effects as an important aspect of their biological function (Flammer and Rogatsky, 2011; Hong and Tontonoz, 2008; Saijo et al., 2009). Investigation of the molecular basis for these actions has revealed diverse points of regulation that can be selectively targeted to achieve receptor- and gene-specific control of the inflammatory response. In the cases of PPAR $\gamma$  and LXRs, for example, ligand-dependent SUMOylation of specific residues in their respective ligand-binding domains results in their interaction with NCoR/SMRT complexes on inflammatory promoters, where they inhibit gene activation by interfering with signal-dependent corepressor clearance (Blaschke et al., 2006; Ghisletti et al., 2009; Pascual et al., 2005; Venteclef et al., 2010). Additional mechanisms utilized by the glucocorticoid receptor and the orphan nuclear receptor Nurr1 include inhibition of specific activator/coactivator interactions and gene-specific recruitment of corepressors (Ogawa et al., 2005; Saijo et al., 2009). Investigation of nuclear receptor repression pathways has thus led to insights into molecular mechanisms that both positively and negatively regulate inflammatory responses at a gene-specific level. To further advance our understanding of these mechanisms, we undertook a genome-wide screen in *Drosophila* to identify novel proteins required for nuclear receptor transrepression. This screen led to the identification and characterization of SMYD5, a previously unrecognized component of the NCoR complex that specifically trimethylates histone H4 at lysine 20 (H4K20me3) on a subset of proinflammatory promoters. This serves as a repression mark that is required for LXR repression and is removed, upon TLR4 activation, by PHF2, a member of the Jumonji domain family of lysine demethylases. These findings highlight a role for active histone H4K20 methylation and demethylation as a molecular checkpoint in the regulation of a subset of proinflammatory gene programs.

## RESULTS

### Identification of SMYD5 as a Negative Regulator of Inflammatory Response Genes


Nuclear receptors exert potent anti-inflammatory effects as important aspects of their biological functions, but these mechanisms remain poorly understood. As *Drosophila melanogaster* utilizes a conserved pathway involving signal-dependent activation of the NF- $\kappa$ B homolog Relish (Rel) to initiate inflammatory responses (Lazzaro, 2008; Lemaitre and Hoffmann, 2007) (Figure S1A) and also expresses nuclear receptors (King-Jones and Thummel, 2005), we explored the use of *Drosophila* Schneider (S2) cells as a model to identify genes involved in

nuclear receptor-mediated repression of inflammatory gene expression. A screen of *Drosophila* nuclear receptors identified Tailless (TII) as an orphan nuclear receptor capable of repressing LPS-dependent activation of an *Attacin A* promoter-luciferase reporter gene (Figure S1B). Using this assay, we evaluated a dsRNA library targeting 21,300 genes for silencing RNAs (Boutros et al., 2004) that could reverse TII-repression of *Attacin A* promoter activity. In addition to known suppressors of inflammatory responses, this screen led to the identification of CG3353 (Figure S1C), encoding a 393 amino acid protein containing a SET domain. CG3353 is most closely related to mammalian SMYD5, a member of the SMYD family of SET and MYND domain-containing proteins (Figure 1A). Using an independent dsRNA, we confirmed that knockdown of CG3353 abolished TII transrepression, as well as resulted in an increase in basal expression of the *Attacin A* reporter gene (Figure 1B).

Primary mouse macrophages express all five members of the mammalian SMYD subfamily (Figure S1D), providing a biologically relevant model system to evaluate their potential roles in innate immune responses. Using specific siRNAs to deplete expression of each SMYD family member (Figure S1E), we observed that knockdown of SMYD5 resulted in exaggerated transcriptional responses of a subset of TLR4-responsive genes following stimulation with a purified LPS, Kdo2 lipid A (KLA) (Figures 1C, 1D, and S1F). In contrast, siRNAs directed against the other members of the mammalian SMYD family (Smyd1–4), resulted in suppressed responses (Figures 1C and S1F), indicating distinct molecular functions. Overexpression of wild-type SMYD5 markedly repressed KLA induction of a TNF $\alpha$ -luciferase reporter gene, while overexpression of SMYD5, containing a mutation predicted to abolish methyltransferase activity (H315L; SMYD5-mut), did not (Figure 1E). Knockdown of SMYD5 expression also abolished or compromised the ability of the LXR-specific agonist (GW3965) to inhibit LPS-induced expression of *Ccl4* (Figure 1F) and other LXR-sensitive genes such as *IL1a* (Figure S1G), suggesting a conserved function in *Drosophila* and mammalian cells. In contrast, knockdown of SMYD5 did not alter inhibitory effects of the glucocorticoid receptor (data not shown), suggesting that SMYD5 contributes to receptor-specific repression mechanisms.

### SMYD5 Trimethylates H4K20 at TLR4-Responsive Promoters

Next, we investigated the ability of bacterially expressed SMYD5 to methylate recombinant histones. Although SMYD1–3 have been shown to methylate histone H3 (Albert and Helin, 2010), a GST-SMYD5 fusion protein purified from bacteria exhibited methyltransferase activity on recombinant histone H4, but not histone H3 (Figure 2A). Mutations in the catalytic domain of SMYD5 (H315L or C317A) abrogated this activity (Figure 2A). To date there are only two identified methylation sites on histone H4: arginine at position 3 and lysine at position 20. Since SET domains typically methylate only lysine residues, our data suggested that SMYD5 most likely functions as an H4K20 methyltransferase. Consistent with this, methylation-specific antibodies detected trimethylation of lysine 20 on histone H4 incubated with wild-type, bacterially expressed SMYD5, but no methylation was observed with mutant SMYD5 (Figure 2B).



**Figure 1. SMYD5 is a Negative Regulator of Inflammatory Response Genes and Is Required for LXR Transrepression**

(A) Percent amino acid identity between *Drosophila* CG3353 and mammalian SMYD5 proteins, which share conserved MYND and SET domains. (B) dsRNA knockdown of CG3353 in *Drosophila* S2 cells reverses TII-dependent repression of the *Attacin A* luciferase reporter. Values represent the mean  $\pm$  SEM of three experiments, \* $p < 0.05$ .

(C) Effect of control (Ctl) or *Smyd1-5* siRNAs transfected into thioglycollate-elicited macrophages on KLA-induced expression of *Ccl4* mRNA 4 hr following treatment. Values represent the average of three experiments  $\pm$  SEM, \* $p < 0.05$  as compared to siCtl LPS treatment.

(D) Effect of control (Ctl) or *Smyd5* siRNAs transfected into thioglycollate-elicited macrophages on KLA-induced expression of *Il1a*, *Il1b*, *Ccl4*, *Cxcl10*, and *Tnf* mRNAs 4 hr following treatment. Values represent the average of three experiments  $\pm$  SEM, \* $p < 0.05$ .


(E) Effect of overexpression of WT SMYD5 and mutant (H315L) SMYD5 on KLA induction of the *Tnf* promoter. Values represent the mean  $\pm$  SEM of three experiments, \* $p < 0.05$ .

(F) Effect of SMYD5 knockdown on LXR repression of *Ccl4* mRNA in thioglycollate-elicited macrophages treated with GW3965 for 1 hr followed by 4 hr of KLA treatment. Values represent the average of three experiments  $\pm$  SEM, \* $p < 0.05$  relative to siCtl, KLA-treated sample.

Immunoprecipitates of FLAG-tagged wild-type and mutant SMYD5 expressed in mammalian cells also catalyzed trimethylation of H4K20, while no methylation was observed for H3K4me3, H3K9me3, H3K27me3, or H3K36me3 (Figure 2C). The specificity of the H4K20me3 antibody was confirmed using recombinant H4 proteins chemically mono-, di-, or trimethylated at position 20 (Figure S2A).

To determine whether H4K20 is the only residue in histone H4 modified by SMYD5 methylation, we performed methyltransferase reactions for SET8, a SET domain protein that monomethylates H4K20, or SMYD5, using recombinant unmodified histone H4 or recombinant H4 chemically trimethylated at position 20. SMYD5 and SET8 methylated unmodified histone H4, but not chemically trimethylated H4K20 (Figure 2D). In addition, FLAG-tagged SMYD5 and SET8 methylated recombinant histone H4, but not recombinant histone H4 in which lysine 20 was mutated to alanine (H4K20A, Figure S2B). Collectively, these results indicate that SMYD5 functions to catalyze trimethylation of H4K20.

(Barski et al., 2007; Gonzalo et al., 2005; Mikkelsen et al., 2007). Therefore, we hypothesized that SMYD5 functions as a corepressor of inflammatory genes by catalyzing local trimethylation of H4K20. Consistent with this, chromatin immunoprecipitation (ChIP) experiments detected H4K20me3 near the transcriptional start sites of several proinflammatory promoters, exemplified by the *Tnf* (Figure 2E) and *Cxcl10* (Figure 2F) promoters, under basal conditions. Notably, there was a marked reduction in this mark upon stimulation of TLR4 with KLA (Figures 2E and 2F). Furthermore, knockdown of SMYD5 by siRNA treatment resulted in an almost complete loss of H4K20me3 on the *Tnf* and *Cxcl10* promoters (Figures 2E and 2F), while not affecting total H4 occupancy (Figures S2C and S2D). *Smyd5* siRNA did not reduce mRNA expression of the previously identified H4K20 methyltransferases: *Set8*, *Nsd1*, *Suv420H1*, or *Suv420H2*, indicating that the observed loss of H4K20me3 on proinflammatory promoters is most likely dependent on reduced SMYD5 expression (Figure S2E). Furthermore, siRNA mediated



**Figure 2. SMYD5 Trimethylates H4K20 at TLR4-Responsive Promoters**

(A) Methyltransferase activity of SMYD5, and two SET domain mutants (H315L and C317A), on recombinant histones H3 and H4. GST-Purified SMYD5 proteins were incubated with radiolabeled H<sup>3</sup>-SAM and recombinant histones for 4 hr. Activity was measured as CPM/μg histone. Values represent the average of three experiments ± SEM, \*p < 0.05.

(B) GST-SMYD5 and GST-SMYD5 (H315L, C317A) were incubated with recombinant histone H4. Reaction products were detected by immunoblotting with antibodies that recognize di- or tri-H4K20-methylation.

(C) Immunoprecipitated FLAG-SMYD5 and FLAG-SMYD5 (H315L/C317A) were incubated with recombinant histone H3 or H4 in the presence or absence of S-adenosylmethionine (SAM). Reaction products were detected by immunoblotting with indicated antibodies.

(D) FLAG-SMYD5 or recombinant SET8 were incubated with recombinant histone H4 or a chemically modified histone H4 trimethylated at K20. Activity was measured as CPM/μg histone. Values represent the average of three experiments ± SEM, \*p < 0.05 relative to Empty, unmodified H4 treatments.

(E) Chromatin immunoprecipitation assays assessing the H4K20me3 levels on the *Tnf* promoter after treatment of thioglycollate-elicited macrophages with *siCtl* and *siSmyd5* for 48 hr, followed by 2 hr of Veh or KLA stimulation. Values represent the average of three experiments ± SEM, \*\*, \*p < 0.05.

(F) Chromatin immunoprecipitation assays assessing H4K20me3 levels on the *Cxcl10* promoter before and after KLA stimulation for 2 hr. Values represent the average of three experiments ± SEM, \*p < 0.05.


knockdown of the previously identified H4K20me3 methyltransferases SUV420H1 and SUV420H2 did not alter the H4K20me3 status of the *Tnf* (Figure S2F) or *Cxcl10* (Figure S2G) promoters, and knockdown of SUV420H1 or SUV420H2 reduced, rather than enhanced, KLA-dependent activation of *Tnf* (Figure S2H) or *Cxcl10* mRNA (Figure S2I).

### SMYD5 Is a Component of NCoR Corepressor Complexes

Next, we sought to determine whether SMYD5 resides at inflammatory promoters. Since available antibodies were not suitable for ChIP, we implemented a strategy to biotin-tag SMYD5 in RAW 264.7 macrophages (de Boer et al., 2003). A biotin ligase recognition peptide (BLRP) was fused to the amino terminus of SMYD5 and the resulting protein was coexpressed with the

of inflammatory signals, whereas activation of TLR4 by KLA resulted in the dismissal of SMYD5 from these promoters (Figures 3A and 3B).

The amino terminus of SMYD5 contains a MYND domain, which was previously demonstrated to mediate interactions with corepressor proteins including NCoR and SMRT (Liu et al., 2007). Gel filtration analysis of nuclear extracts from HEK293 cells transfected with GFP-SMYD5 and FLAG-NCoR indicated that GFP-SMYD5 eluted with NCoR and HDAC3 in high molecular weight fractions (Figure 3C). Coimmunoprecipitation analysis in 293T cells detected interaction between full-length SMYD5 and NCoR (Figure 3D). This interaction suggested that the NCoR complex could possess H4K20 histone methyltransferase activity. To test this hypothesis, we expressed FLAG-NCoR in 293T cells, purified the complex using an anti-FLAG affinity



**Figure 3. The NCoR Complex Directs SMYD5 to TLR4-Responsive Promoters**

(A) Chromatin immunoprecipitation assays assessing BLRP-SMYD5 occupancy of the *Tnf* promoter before and after KLA stimulation for 2 hr. Values represent the average of three experiments  $\pm$  SEM,  $^*p < 0.05$ .  
 (B) Chromatin immunoprecipitation assays assessing BLRP-SMYD5 occupancy of the *Cxcl10* promoter before and after KLA stimulation for 2 hr. Values represent the average of three experiments  $\pm$  SEM,  $^*p < 0.05$ .  
 (C) Gel filtration experiments for HEK293 cell lysates transfected with FLAG-NCoR and GFP-SMYD5. The lysates were then separated using SDS-PAGE analysis and immunoblotted with anti-NCoR, HDAC3, and GFP antibodies.  
 (D) Coimmunoprecipitation of BLRP-SMYD5 with FLAG-NCoR in BLRP-SMYD5 RAW 264.7 cells.  
 (E) Methyltransferase reaction for SMYD5 and NCoR. HEK293 cells were transfected with vectors containing FLAG-Empty, FLAG-SMYD5, or FLAG-NCoR and immunoprecipitated using FLAG M2 beads. Lysates were subjected to a methyltransferase reaction and immunoblotted using a specific H4K20me3 antibody.  
 (F) Chromatin immunoprecipitation assays assessing the H4K20me3 levels on the *Tnf* and *Cxcl10* promoters in WT and NCoR<sup>-/-</sup> bone marrow-derived macrophages. Values represent the average of three experiments  $\pm$  SEM,  $^*p < 0.05$ .  
 (G) Chromatin immunoprecipitation assays assessing the BLRP-SMYD5 recruitment to the *Ccl4*, *Tnf*, and *Il6* promoters following *shCt1* and *shNCoR* transfected Raw-BLRP-SMYD5 cells. Values represent the average of three experiments  $\pm$  SEM,  $^*p < 0.05$ .

matrix, and subjected the precipitate to histone methyltransferase assays using recombinant histone H4 as a substrate. Interestingly, both the SMYD5 and NCoR complexes methylated

H4K20me3 (Figure 3E). Genetic deletion of NCoR in bone marrow-derived macrophages resulted in the loss of H4K20me3 on proinflammatory promoters, including *Tnf* and *Cxcl10*, as compared to WT macrophages (Figure 3F). Furthermore, SMYD5 occupancy on proinflammatory promoters was significantly reduced in BLRP-SMYD5 expressing RAW 264.7 cells following shRNA-mediated knockdown of NCoR (Figure 3G).

### PHF2 Is Required for Removal of H4K20me3 and Activation of TLR4-Responsive Promoters


The observation that the H4K20me3 mark at inducible promoters was markedly reduced shortly after TLR4 ligation suggested the possibility of signal-dependent demethylation. Although no H4K20me3 demethylases have been described, recent studies have shown that PHF2 demethylates H4K20me1 (Liu et al., 2010; Qi et al., 2010), raising the possibility that two closely related proteins previously shown to possess demethylase activity, PHF2 (Wen et al., 2010) and KIAA1718 (Horton et al., 2010), may target H4K20me3 on inflammatory gene promoters. Knockdown of Phf2, but not Kiaa1718 or Phf8 (Figures S3A–S3C), significantly diminished KLA induction of the SMYD5-sensitive *Tnf* and *Cxcl10* genes (Figures 4A and 4B).

To investigate PHF2 enzymatic function, wild-type PHF2 and a mutant PHF2-A/A (H248A, D250A), which alters conserved residues required for iron binding in the active site of the Jumonji domain, were purified from bacteria cell extracts and subjected to an in vitro histone demethylation assay using purified core histones and mononucleosomes. PHF2 effectively removed H4K20me3 methylation on mononucleosomes, while the mutant PHF2-A/A was much less active (Figures 4C and 4D). PHF2 also demethylated H3K9me1 on mononucleosomes as well as on core histones, consistent with recent observations (Wen et al., 2010) (Figure 4C).

ChIP experiments were next performed to investigate whether PHF2 was involved in the removal of the H4K20me3 mark at TLR4-responsive promoters. Primary bone marrow-derived macrophages were transfected with control siRNAs or siRNAs directed against *Phf2*, and the cells were then treated with vehicle or KLA for 4 hr. The H4K20me3 mark was depleted from the *Tnf* and *Cxcl10* promoters in siCt1-treated cells upon KLA treatment, as previously seen, but not in PHF2 knockdown macrophages (Figures 4E and 4F). Next, we sought to determine whether PHF2 is recruited to promoters exhibiting loss of H4K20me3 in response to LPS signaling. Because available antibodies were not suitable for the ChIP studies of PHF2, we implemented the above described biotin-tagging strategy to perform ChIP studies of PHF2 in RAW 264.7 macrophages. These studies demonstrated that PHF2 is recruited to the proinflammatory promoters *Tnf* and *Cxcl10*, but not  $\beta$ -actin, only upon activation of TLR4 signaling, correlating with the removal of H4K20me3 (Figure 4G).

### Exchange of SMYD5 and PHF2 Controls the Regulation of TLR4-Dependent Gene Expression

To globally evaluate the roles of PHF2 and SMYD5 in the regulation of TLR4-dependent gene activation, we performed RNA-sequencing of polyA RNA isolated from primary macrophages following transfections with *Ctrl*, *Smyd5*, or *Phf2* siRNAs for



**Figure 4. PHF2 Demethylates H4K20me3 and Is Required for TLR4-Dependent Gene Activation**

(A) Quantitative real-time PCR for *Tnf* mRNA isolated from thioglycollate-elicited macrophages treated with siRNA for Control, *Phf2*, *Phf8*, and *Kiaa1718* and subsequently treated with KLA for 4 hr. Values represent the average of three experiments  $\pm$  SEM, \* $p < 0.05$ .

(B) Quantitative real-time PCR for *Cxcl10* mRNA isolated from thioglycollate-elicited macrophages treated with siRNA for Control, *Phf2*, *Phf8*, and *Kiaa1718* and subsequently treated with KLA for 4 hr. Values represent the average of three experiments  $\pm$  SEM, \* $p < 0.05$ .

(C) Histone demethylase assay for PHF2 and a PHF2 mutant performed on core histone and mononucleosomes. His-PHF2(wt) or His-PHF2(mut) (H248A, D250A) were purified and incubated with core histones or mononucleosomes in histone demethylation buffer for 4 hr. Demethylation activity was evaluated by immunoblotting with specific antibodies.

(D) Quantification of H4K20me3 intensities from Figure 3C normalized to H4 signal.


(E) Chromatin immunoprecipitation assays assessing the occupancy of H4K20me3 on the *Tnf* promoter in thioglycollate-elicited macrophages treated with Veh or KLA stimulation for 2 hr. Values represent the average of three experiments  $\pm$  SEM, \* $p < 0.05$ .

(F) Chromatin immunoprecipitation assays assessing the occupancy of H4K20me3 on the *Cxcl10* promoter in thioglycollate-elicited macrophages treated with Veh or KLA stimulation for 2 hr. Values represent the average of three experiments  $\pm$  SEM, \* $p < 0.05$ .

(G) Chromatin immunoprecipitation assays assessing the recruitment of BLRP-PHF2 to the  $\beta$ -actin, *Cxcl10*, and *Tnf* promoters upon KLA stimulation for 2 hr. Values represent the average of three experiments  $\pm$  SEM, \* $p < 0.05$ .

48 hr and subsequently treated with KLA for 4 hr (Table S1S). Of the 419 mRNAs that were significantly ( $FDR < 0.00001$ ) stimulated greater than 2-fold by KLA treatment (Figure 5A), knockdown of PHF2 blunted the response of 51% (214 of 419) of these mRNAs. PHF2-dependent mRNAs were significantly enriched for functional annotations linked to cytokine activity, chemokine receptor binding and chemotaxis (Figure S4A). Conversely, knockdown of SMYD5 resulted in significantly exaggerated KLA responses for 63 of these mRNAs, which were significantly enriched for functional annotations linked to immune, defense, and inflammatory responses (Figure S4B). Notably, 68% (43 of

63) of mRNAs that were hyperinduced by knockdown of SMYD5 required PHF2 for activation ( $p < 0.005$ ). However, the observation that the PHF2-dependent subset of TLR4-responsive genes was substantially larger than the subset of genes hyperinduced following SMYD5 knockdown raised the question of whether the actual set of H4K20me3-marked promoters are more broadly distributed than that defined by hyperinduction. Consistent with this possibility, ChIP experiments indicated that H4K20me3 was present on the promoters of 10/10 PHF2-dependent KLA target genes evaluated, regardless of whether they were hyperactivated following SMYD5 depletion (Figure 5B). These results suggest that SMYD5 and PHF2 function in a reciprocal manner to regulate the transcriptional response to KLA. Consistent with this possibility, knockdown of *Smyd5* mRNA, which depletes SMYD5-dependent promoters of H4K20me3, circumvents the



**Figure 5. Exchange of SMYD5 and PHF2 Controls the Regulation of TLR4-Dependent Gene Expression**

(A) Venn diagram comparing the overlap of LPS genes that are blocked by *siPhf2* treatment, enhanced by *siSmyd5* treatment, or unaffected by either treatment identified by polyA mRNA sequencing.

(B) Chromatin immunoprecipitation assays for H4K20me3 on SMYD5-dependent and SMYD5-independent promoters in thioglycollate-elicited macrophages. Values represent the average of three experiments ± SEM, \*p < 0.05.

(C) Quantitative real-time PCR for *Tnf* and *Cxcl10* mRNAs isolated from thioglycollate-elicited macrophages treated with siRNA for *Control*, *Phf2*, *Smyd5*, or a combination of *Smyd5* and *Phf2* siRNA, and subsequently treated with KLA for 4 hr. Values represent the average of three experiments ± SEM, \*p < 0.05.

(D) De novo motif analysis of PHF2-dependent promoters.

(E) Coimmunoprecipitation assay for transfected FLAG-PHF2 with endogenous p65 in RAW 264.7 cells treated with Veh or KLA for 1 hr.

(F) Chromatin immunoprecipitation assays for BLRP-PHF2 on the *Tnf* and *Cxcl10* promoters in cells transfected with *shCtl* or *sh-p65* for 48 hr and subsequently treated with KLA for 3 hr. Values represent the average of three experiments ± SEM, \*p < 0.05.

(G) Chromatin immunoprecipitation assays for H4K20me3 on the *Ccl4* promoter in thioglycollate-elicited macrophages treated with Veh, KLA, or a combination of GW3965 and KLA. Values represent the average of three experiments ± SEM, \*p < 0.05.

necessity for PHF2 in LPS-dependent gene activation of the *Tnf* and *Cxcl10* mRNAs (Figure 5C).


To gain insights into potential proteins involved in the signal-dependent recruitment of PHF2 to these promoters, we analyzed the promoters of PHF2-dependent target genes and found NF-κB and IRF recognition motifs to be highly enriched (Figure 5D). Therefore, we performed coimmunoprecipitation assays to determine whether PHF2 interacts with p65, a member of the NF-κB family, which is required for activation of a large subset of TLR4-dependent genes (Lim et al., 2007; Medzhitov and Horng, 2009). We detected KLA-dependent interaction between p65 and PHF2 (Figure 5E). Notably, knockdown of p65 resulted in diminished recruitment of PHF2 to both the *Cxcl10* and *Tnf* promoters, demonstrating that p65 is required for full recruitment of PHF2 (Figure 5F).

LXRs exert repressive effects on TLR4 target genes by preventing signal-dependent NCoR clearance (Ghisletti et al.,

this possibility, the LXR agonist GW3965 inhibited KLA-induced erasure of H4K20me3 on the *Ccl4* promoter (Figure 5G).

## DISCUSSION

In concert, these findings reveal remarkable specificity in the growing spectrum of biological processes controlled by regulated methylation/demethylation of H4K20, previously established to play roles in cell cycle (H4K20me1) (Liu et al., 2010; Wu et al., 2010), DNA damage repair (H4K20me2) (Pei et al., 2011; Sanders et al., 2004), and Wnt signaling (Li et al., 2011). Here, we provide evidence for an H4K20me3 methylation/demethylation mechanism, catalyzed by SMYD5 and PHF2, respectively, which is required for signal-dependent regulation of a subset of inflammatory response genes. We propose that under resting conditions, SMYD5 is recruited to a subset of TLR4-responsive promoters through its association with NCoR



**Figure 6. Role of H4K20me3 in Regulation of TLR4-Responsive Genes**

Integrated model for SMYD5 and PHF2 regulation of inflammatory gene promoters.

corepressor complexes, where it trimethylates H4K20 (Figure 6). Thus, in addition to mediating repression through the removal of acetylation by associated histone deacetylases, such as HDAC3, NCoR complexes also contribute to repression through the histone methyltransferase activity of associated SMYD5. The basis for the repressive function of H4K20me3 remains to be established; however, recent studies have suggested that H4K20me3 inhibits acetylation of H4K16, resulting in the pausing of RNA polymerase II (Kapoor-Vazirani et al., 2011). Consistent with this, paused Pol II is a characteristic feature of many genes that are rapidly and highly induced by TLR4 signaling (Escoubet-Lozach et al., 2011; Hargreaves et al., 2009).

The present studies expand on the emerging recognition of distinct biochemical activities and biological functions of members of the SMYD family of histone methyltransferases. SMYD1–3 have been reported to mediate trimethylation of histone H3K4, and reduced deposition of this activation mark could potentially account for why siRNA knockdown of *Smyd1*, 2, and 3 resulted in reduced responses to TLR4 activation (Albert and Helin, 2010). It will thus be of interest to define the molecular mechanisms that dictate distinct substrate specificities. The preference of SMYD5 for H4K20 appears to be an intrinsic property of the enzyme based on the observed activity of the purified GST-SMYD5 expressed in bacteria. In addition to SMYD5 and SUV420H1/2, SMYD3 has also recently been reported to trimethylate H4K20 (Foreman et al., 2011). However, the observations that knockdown of each of these enzymes resulted in reduced activation of TLR4 target genes, and that the presence of the H4K20me3 mark at TLR4-responsive promoters was largely dependent on SMYD5, strongly argue that SMYD3 and SUV420H1/2 primarily act on different substrates and/or are directed to H4K20 at different genomic locations by alternative protein complexes. Of note, we find that knockdown of SUV420H1/2 results in significant reduction in the expression of Phf2, which could at least partly explain the effect of their knockdown on TLR4-dependent gene activation (J.D.S., unpublished data).

We were initially led to SMYD5 by an RNAi screen that identified the *Drosophila* homolog CG3353 as being required for Tailless repression of LPS signaling. The observation that SMYD5 is required for transrepression activities of LXRs on at

least a subset of its target genes suggests evolutionarily conserved molecular functions of SMYD5 that link nuclear receptor signaling to antagonism of innate immune responses. The finding that SMYD5 is required for LXR-dependent, but not glucocorticoid receptor-dependent, antagonism of TLR4-induced gene expression is also consistent with prior observations that LXR, but not glucocorticoid receptor, transrepression requires NCoR (Blaschke et al., 2006; Ghisletti et al., 2009; Venticlef et al., 2010; Ogawa et al., 2005).

We further show that TLR4-induced removal of H4K20me3 is catalyzed by PHF2, which is delivered to responsive promoters by the p65 subunit of NF-κB (Figure 6). The observation that more genes require PHF2 for optimal responses to TLR4 signaling than are hyperinduced in the setting of SMYD5 knockdown has several potential explanations. One is that while removal of H4K20me3 may be required for signal-dependent activation, lack of this mark may not be sufficient for hyperactivation, suggesting that it serves a repression “checkpoint” function. Consistent with this possibility, many PHF2-dependent promoters that were not hyperresponsive in response to SMYD5 knockdown were nevertheless marked by H4K20me3 under basal conditions. It is also possible that PHF2 targets additional methylated residues, such as H3K9me1, on proinflammatory promoters that are deposited by other histone methyltransferases. It is of interest that PHF2 was capable of catalyzing removal of H3K9me1, but not H4K20me3, on core histones. We speculate that the requirement for a nucleosome substrate for demethylation of H4K20me3 may be due to the ability of the plant homeodomain in PHF2 to recognize H3K4me3 (Wen et al., 2010). This interaction may anchor the enzyme in the correct conformation or allosterically regulate substrate specificity, as previously suggested for H4K20me1 demethylation by PHF8 (Liu et al., 2010). As our findings suggest that demethylation of H4K20me3 is a critical step in gene activation of inflammatory response genes, such as *Tnf*, modulation of the expression or activity of PHF2 may provide novel approaches for treatment of inflammatory diseases.

## EXPERIMENTAL PROCEDURES

### *Drosophila* RNAi Screen

Schneider S2 cells were plated in 6 cm dishes and transfected with 1.5 μg *Attacin A* reporter and 1.5 μg Tailless expression vector using Effectene (QIAGEN) transfection reagent according to the manufacturer’s guidelines. The cells were then plated in 384-well plates containing 250 ng of dsRNA and incubated for 48 hr. Cells were then treated with 100 ng/ml LPS for 8 hr and lysed, and luciferase activity was monitored using a luminometer.

### Gel Filtration

HEK293 cells were transfected with FLAG-tagged NCoR WT and GFP-SMYD5 WT using Lipofectamine 2000. Forty-eight hours post transfection, HEK293 cultured cells were harvested and washed twice with ice-cold PBS, and cytoplasmic extracts were prepared using a low-salt buffer (10 mM HEPES [pH 7.9], 50 mM NaCl, 1 mM DTT). Nuclei were spun down and extracted in 20 mM HEPES (pH 7.9), 20% glycerol, 350 mM NaCl, 0.2 EDTA, 1 mM DTT, and protease inhibitors. A 1 mg sample was then applied onto a 250 ml column of Sephacryl S-300 (Amersham Biosciences, Inc.) previously equilibrated with gel filtration buffer (20 mM HEPES [pH 7.9], 120 mM NaCl, 0.2 mM EDTA, 1 mM DTT) and 60 1 ml fractions were collected. Fractions were analyzed by western blot.



### Cell Culture and Western Blots

Raw 264.7 cells were grown in DMEM supplemented with 10% FBS and penicillin/streptomycin. Raw 264.7 cells were transfected using Superfect (QIAGEN) according to the manufacturer's guidelines and as previously published (Ghisletti et al., 2009). Primary thioglycollate-elicited macrophages and bone marrow-derived macrophage cells were generated as previously described (Ogawa et al., 2005). BLRP stables for SMYD5 and PHF2 were generated as previously reported (Heinz et al., 2010). LPS (Sigma) and KLA (Avanti) were used at a concentration of 100 ng/ml unless otherwise stated. The shRNA constructs were constructed using pSilencer 3.0 H1 (Ambion). Antibodies used are as follows: FLAG (Sigma, F3165), GFP (Abcam, Ab290), p65 (Santa Cruz, sc-8008), BLRP (Genscript, A00674), HDAC3 (Santa Cruz, sc-11417), H4 (Active Motif, 39269), H4K20me1 (Abcam, Ab9051), H4K20me2 (Abcam, Ab9052), H4K20me3 (Abcam, Ab9053), H3K9me1 (Abcam, Ab8896), H3K9me2 (Abcam, Ab1220), H3K9me3 (Abcam, Ab8898), H3K27me2 (Abcam, Ab24684), H3K27me3 (Millipore, 07-449) H3K4me3 (Abcam, AB8580), H3K36me3 (Abcam, Ab9050), H3 (Active Motif, 39163), and His (Santa Cruz, SC-803). Chemically methylated H4 proteins were purchased from Active Motif.

### RNA Isolation, Quantification, and Sequencing

RNA was purified using RNeasy Mini Kit (QIAGEN) and enriched for Poly(A)-RNA with MicroPoly(A) Purist Kit (Ambion, Austin, TX). For quantification, cDNA was generated using the Superscript kit (Invitrogen), and quantitative real-time PCR was performed using gene-specific primers. For sequencing, RNA was treated with TURBO DNase (Ambion), fragmented using RNA Fragmentation Reagents (Ambion) and purified by a P-30 column (Bio-Rad, Hercules, CA, USA). Fragmented RNA was dephosphorylated with Antarctic phosphatase (New England Biolabs, Ipswich, MA) followed by heat-inactivation and overnight precipitation. Poly(A)-tailing and cDNA synthesis was performed as previously described (Ingolia et al., 2009). However, for reverse transcription, oligos with custom barcodes (underlined) were used: 5'-Phos CA/TG/AC/GT GATCGTCGGACTGTAGAACTCT/idSp/CAAGCAGAAGACGGCA TACGATTTTTTTTTTTTTTTTTTTTTVN-3'. After cDNA synthesis, exonuclease was used to catalyze the removal of excess oligo. Enzyme was inactivated and RNA hydrolyzed by alkaline treatment (100 mM NaOH) and heat (25 min, 95°C). The cDNA fragments of 50–150 nucleotides were purified on a denaturing Novex 10% polyacrylamide TBE-urea gel (Invitrogen, Carlsbad, CA). The recovered cDNA was circularized, linearized, amplified for 15 cycles, and gel purified as previously described (Ingolia et al., 2009). The library was sequenced on the Illumina Genome Analyzer 2 according to the manufacturer's instructions. Gene Ontology analysis for regulated genes was performed as previously described (Sullivan et al., 2011). Promoters of regulated genes were analyzed for enriched motifs as previously described (Heinz et al., 2010).

### siRNA Transfections

Transfections using 5 nM siRNA were performed using lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. SMART siRNA pools for examined genes were purchased from Dharmacon.

### Histone Methylation Assays

Purification of GST-SMYD5, GST-SMYD5 mutants, His-H4, and His-H4K20A was performed using standard molecular biology techniques. FLAG-NCoR and FLAG-SMYD5 complexes were obtained from transiently transfected HEK293 cells and immunoprecipitated with FLAG M2 beads (Sigma). Purified proteins were incubated with 2  $\mu$ g recombinant histone H4 (NEB), or chemically methylated histone H4 (Active Motif), or His-tagged H4 or H4K20A, 1 ml [<sup>3</sup>H] adenosyl-L-methionine (PerkinElmer) or 0.3 mM SAM (NEB), in histone methylation buffer (50 mM Tris [pH 9.0], 0.5 mM DTT) overnight at 30°C. For the radiolabeled experiments, the reactions were then transferred to P81 paper and washed three times with 10% trichloroacetic acid for 15 min, followed by a wash with 95% ethanol, and analyzed using a scintillation counter. Otherwise, samples were separated using SDS-PAGE and immunoblotted using specific antibodies.

### ChIP Assay

ChIP assays were performed as previously described (Heinz et al., 2010). The antibodies used in these studies are as follows: IgG (Santa Cruz Biotech-

nology), H4K20me3 (Abcam, Ab9053). SMYD5 and PHF2 ChIPs were performed using streptavidin immunoprecipitations from Raw 264.7 cells stably expressing SMYD5 or PHF2 tagged with a biotin recognition peptide as described previously (Heinz et al., 2010).

### Demethylation Assay

His-PHF2 and His-PHF2 (H248A, D250A) were purified from bacterial extracts using standard protocols and were incubated with 2  $\mu$ g mononucleosomes in DeMTase buffer (20 mM Tris-HCl [pH 7.5], 150 mM NaCl, 50 mM [NH<sub>4</sub>]<sub>2</sub>Fe[SO<sub>4</sub>]<sub>2</sub>) for 2–5 hr at 37°C. Samples were separated using SDS-PAGE and immunoblotted using specific antibodies. Quantification of western blots were performed using ImageJ software (Abramoff et al., 2004).

### Coimmunoprecipitations

FLAG-PHF2 or FLAG-NCoR and BLRP-SMYD5 were transfected into 293T cells. Cells were lysed and sonicated in lysis buffer (10 mM Tris [pH 8.0], 420 mM NaCl, 0.5% NP40, 1mM EDTA, Protease Inhibitors [Roche]). Samples were then diluted with dilution buffer (10 mM Tris [pH 8.0], 0.5% NP40, 1.0 mM EDTA) and protein complexes were purified using FLAG M2 beads (Sigma). Beads were washed with dilution buffer five times and eluted with 3X FLAG peptide. Lysates were then separated using SDS-PAGE and interactions were detected using western blot analysis.

### ACCESSION NUMBERS

Sequencing data are available in the Gene Expression Omnibus database (<http://www.ncbi.nlm.nih.gov/geo>) under the accession number GSE39113.

### SUPPLEMENTAL INFORMATION

Supplemental Information includes four figures and one table and can be found with this article online at <http://dx.doi.org/10.1016/j.molcel.2012.07.020>.

### ACKNOWLEDGMENTS

We thank Dawn Zhang and David Gosselin for critical comments and Lynn Bautista for assistance with preparation of the manuscript. These studies were supported by P01-HC088093, P30 DK063491, and CA52599 to C.K.G. M.G.R. is supported by the Howard Hughes Medical Institute. J.S. was supported by T32 CA009523. M.U.K. was supported by a Fondation Leducq Career Development award and a Sigrid Jesélius fellowship.

Received: February 29, 2012

Revised: June 21, 2012

Accepted: July 19, 2012

Published online: August 23, 2012

### REFERENCES

- Abramoff, M.D., Magalhães, P.J., and Ram, S.J. (2004). Image Processing with ImageJ. *Biophotonics International* 11, 36–42.
- Albert, M., and Helin, K. (2010). Histone methyltransferases in cancer. *Semin. Cell Dev. Biol.* 21, 209–220.
- Barski, A., Cuddapah, S., Cui, K., Roh, T.Y., Schones, D.E., Wang, Z., Wei, G., Chepelev, I., and Zhao, K. (2007). High-resolution profiling of histone methylations in the human genome. *Cell* 129, 823–837.
- Blaschke, F., Takata, Y., Caglayan, E., Collins, A., Tontonoz, P., Hsueh, W.A., and Tangirala, R.K. (2006). A nuclear receptor corepressor-dependent pathway mediates suppression of cytokine-induced C-reactive protein gene expression by liver X receptor. *Circ. Res.* 99, e88–e99.
- Boutros, M., Kiger, A.A., Armknecht, S., Kerr, K., Hild, M., Koch, B., Haas, S.A., Paro, R., and Perrimon, N.; Heidelberg Fly Array Consortium. (2004). Genome-wide RNAi analysis of growth and viability in *Drosophila* cells. *Science* 303, 832–835.

- de Boer, E., Rodriguez, P., Bonte, E., Krijgsvelde, J., Katsantoni, E., Heck, A., Grosveld, F., and Strouboulis, J. (2003). Efficient biotinylation and single-step purification of tagged transcription factors in mammalian cells and transgenic mice. *Proc. Natl. Acad. Sci. USA* *100*, 7480–7485.
- Escoubet-Lozach, L., Benner, C., Kaikkonen, M.U., Lozach, J., Heinz, S., Spann, N.J., Crotti, A., Stender, J., Ghisletti, S., Reichart, D., et al. (2011). Mechanisms establishing TLR4-responsive activation states of inflammatory response genes. *PLoS Genet.* *7*, e1002401.
- Flammer, J.R., and Rogatsky, I. (2011). Minireview: Glucocorticoids in autoimmunity: unexpected targets and mechanisms. *Mol. Endocrinol.* *25*, 1075–1086.
- Foreman, K.W., Brown, M., Park, F., Erntage, S., Harriss, J., Das, C., Zhu, L., Crew, A., Arnold, L., Shaaban, S., and Tucker, P. (2011). Structural and functional profiling of the human histone methyltransferase SMYD3. *PLoS ONE* *6*, e22290.
- Ghisletti, S., Huang, W., Jepsen, K., Benner, C., Hardiman, G., Rosenfeld, M.G., and Glass, C.K. (2009). Cooperative NCoR/SMRT interactions establish a corepressor-based strategy for integration of inflammatory and anti-inflammatory signaling pathways. *Genes Dev.* *23*, 681–693.
- Glass, C.K., Saijo, K., Winner, B., Marchetto, M.C., and Gage, F.H. (2010). Mechanisms underlying inflammation in neurodegeneration. *Cell* *140*, 918–934.
- Gonzalo, S., García-Cao, M., Fraga, M.F., Schotta, G., Peters, A.H., Cotter, S.E., Eguía, R., Dean, D.C., Esteller, M., Jenuwein, T., and Blasco, M.A. (2005). Role of the RB1 family in stabilizing histone methylation at constitutive heterochromatin. *Nat. Cell Biol.* *7*, 420–428.
- Hargreaves, D.C., Horng, T., and Medzhitov, R. (2009). Control of inducible gene expression by signal-dependent transcriptional elongation. *Cell* *138*, 129–145.
- Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* *38*, 576–589.
- Hoberg, J.E., Popko, A.E., Ramsey, C.S., and Mayo, M.W. (2006). IkappaB kinase alpha-mediated derepression of SMRT potentiates acetylation of RelA/p65 by p300. *Mol. Cell Biol.* *26*, 457–471.
- Hong, C., and Tontonoz, P. (2008). Coordination of inflammation and metabolism by PPAR and LXR nuclear receptors. *Curr. Opin. Genet. Dev.* *18*, 461–467.
- Horton, J.R., Upadhyay, A.K., Qi, H.H., Zhang, X., Shi, Y., and Cheng, X. (2010). Enzymatic and structural insights for substrate specificity of a family of jumonji histone lysine demethylases. *Nat. Struct. Mol. Biol.* *17*, 38–43.
- Hotamisligil, G.S. (2006). Inflammation and metabolic disorders. *Nature* *444*, 860–867.
- Huang, W., Ghisletti, S., Perissi, V., Rosenfeld, M.G., and Glass, C.K. (2009). Transcriptional integration of TLR2 and TLR4 signaling at the NCoR derepression checkpoint. *Mol. Cell* *35*, 48–57.
- Huang, W., Ghisletti, S., Saijo, K., Gandhi, M., Aouadi, M., Tesz, G.J., Zhang, D.X., Yao, J., Czech, M.P., Goode, B.L., et al. (2011). Coronin 2A mediates actin-dependent de-repression of inflammatory response genes. *Nature* *470*, 414–418.
- Ingolia, N.T., Ghaemmaghami, S., Newman, J.R., and Weissman, J.S. (2009). Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. *Science* *324*, 218–223.
- Kapoor-Vazirani, P., Kagey, J.D., and Vertino, P.M. (2011). SUV420H2-mediated H4K20 trimethylation enforces RNA polymerase II promoter-proximal pausing by blocking hMOF-dependent H4K16 acetylation. *Mol. Cell Biol.* *31*, 1594–1609.
- Karin, M., Lawrence, T., and Nizet, V. (2006). Innate immunity gone awry: linking microbial infections to chronic inflammation and cancer. *Cell* *124*, 823–835.
- Kawai, T., and Akira, S. (2010). The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat. Immunol.* *11*, 373–384.
- King-Jones, K., and Thummel, C.S. (2005). Nuclear receptors—a perspective from *Drosophila*. *Nat. Rev. Genet.* *6*, 311–323.
- Kourmouli, N., Sun, Y.M., van der Sar, S., Singh, P.B., and Brown, J.P. (2005). Epigenetic regulation of mammalian pericentric heterochromatin in vivo by HP1. *Biochem. Biophys. Res. Commun.* *337*, 901–907.
- Lazzaro, B.P. (2008). Natural selection on the *Drosophila* antimicrobial immune system. *Curr. Opin. Microbiol.* *11*, 284–289.
- Lemaître, B., and Hoffmann, J. (2007). The host defense of *Drosophila melanogaster*. *Annu. Rev. Immunol.* *25*, 697–743.
- Li, Z., Nie, F., Wang, S., and Li, L. (2011). Histone H4 Lys 20 monomethylation by histone methylase SET8 mediates Wnt target gene activation. *Proc. Natl. Acad. Sci. USA* *108*, 3116–3123.
- Lim, C.A., Yao, F., Wong, J.J., George, J., Xu, H., Chiu, K.P., Sung, W.K., Lipovich, L., Vega, V.B., Chen, J., et al. (2007). Genome-wide mapping of RELA(p65) binding identifies E2F1 as a transcriptional activator recruited by NF-kappaB upon TLR4 activation. *Mol. Cell* *27*, 622–635.
- Liu, Y., Chen, W., Gaudet, J., Cheney, M.D., Roudaia, L., Cierpicki, T., Klet, R.C., Hartman, K., Laue, T.M., Speck, N.A., and Bushweller, J.H. (2007). Structural basis for recognition of SMRT/N-CoR by the MYND domain and its contribution to AML1/ETO's activity. *Cancer Cell* *11*, 483–497.
- Liu, W., Tanasa, B., Tyurina, O.V., Zhou, T.Y., Gassmann, R., Liu, W.T., Ohgi, K.A., Benner, C., Garcia-Bassets, I., Aggarwal, A.K., et al. (2010). PHF8 mediates histone H4 lysine 20 demethylation events involved in cell cycle progression. *Nature* *466*, 508–512.
- Martinelli, C., and Reichhart, J.M. (2005). Evolution and integration of innate immune systems from fruit flies to man: lessons and questions. *J. Endotoxin Res.* *11*, 243–248.
- Medzhitov, R., and Horng, T. (2009). Transcriptional control of the inflammatory response. *Nat. Rev. Immunol.* *9*, 692–703.
- Mikkelsen, T.S., Ku, M., Jaffe, D.B., Issac, B., Lieberman, E., Giannoukos, G., Alvarez, P., Brockman, W., Kim, T.K., Koche, R.P., et al. (2007). Genome-wide maps of chromatin state in pluripotent and lineage-committed cells. *Nature* *448*, 553–560.
- Ogawa, S., Lozach, J., Jepsen, K., Sawka-Verhelle, D., Perissi, V., Sasik, R., Rose, D.W., Johnson, R.S., Rosenfeld, M.G., and Glass, C.K. (2004). A nuclear receptor corepressor transcriptional checkpoint controlling activator protein 1-dependent gene networks required for macrophage activation. *Proc. Natl. Acad. Sci. USA* *101*, 14461–14466.
- Ogawa, S., Lozach, J., Benner, C., Pascual, G., Tangirala, R.K., Westin, S., Hoffmann, A., Subramaniam, S., David, M., Rosenfeld, M.G., and Glass, C.K. (2005). Molecular determinants of crosstalk between nuclear receptors and toll-like receptors. *Cell* *122*, 707–721.
- Pascual, G., Fong, A.L., Ogawa, S., Gamlie, A., Li, A.C., Perissi, V., Rose, D.W., Willson, T.M., Rosenfeld, M.G., and Glass, C.K. (2005). A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR-gamma. *Nature* *437*, 759–763.
- Pei, H., Zhang, L., Luo, K., Qin, Y., Chesni, M., Fei, F., Bergsagel, P.L., Wang, L., You, Z., and Lou, Z. (2011). MMSET regulates histone H4K20 methylation and 53BP1 accumulation at DNA damage sites. *Nature* *470*, 124–128.
- Qi, H.H., Sarkissian, M., Hu, G.Q., Wang, Z., Bhattacharjee, A., Gordon, D.B., Gonzales, M., Lan, F., Ongusaha, P.P., Huarte, M., et al. (2010). Histone H4K20/H3K9 demethylase PHF8 regulates zebrafish brain and craniofacial development. *Nature* *466*, 503–507.
- Saijo, K., Winner, B., Carson, C.T., Collier, J.G., Boyer, L., Rosenfeld, M.G., Gage, F.H., and Glass, C.K. (2009). A Nurr1/CoREST pathway in microglia and astrocytes protects dopaminergic neurons from inflammation-induced death. *Cell* *137*, 47–59.
- Sanders, S.L., Portoso, M., Mata, J., Bähler, J., Allshire, R.C., and Kouzarides, T. (2004). Methylation of histone H4 lysine 20 controls recruitment of Crb2 to sites of DNA damage. *Cell* *119*, 603–614.
- Schotta, G., Lachner, M., Sarma, K., Ebert, A., Sengupta, R., Reuter, G., Reinberg, D., and Jenuwein, T. (2004). A silencing pathway to induce H3-K9

and H4-K20 trimethylation at constitutive heterochromatin. *Genes Dev.* **18**, 1251–1262.

Sullivan, A.L., Benner, C., Heinz, S., Huang, W., Xie, L., Miano, J.M., and Glass, C.K. (2011). Serum response factor utilizes distinct promoter- and enhancer-based mechanisms to regulate cytoskeletal gene expression in macrophages. *Mol. Cell Biol.* **31**, 861–875. Published online December 6, 2010. <http://dx.doi.org/10.1128/MCB.00836-10>.

Takeuchi, O., and Akira, S. (2010). Pattern recognition receptors and inflammation. *Cell* **140**, 805–820.

Tedgui, A., and Mallat, Z. (2006). Cytokines in atherosclerosis: pathogenic and regulatory pathways. *Physiol. Rev.* **86**, 515–581.

Venteclef, N., Jakobsson, T., Ehrlund, A., Damdimopoulos, A., Mikkonen, L., Ellis, E., Nilsson, L.M., Parini, P., Jänne, O.A., Gustafsson, J.A., et al. (2010). GPS2-dependent corepressor/SUMO pathways govern anti-inflammatory actions of LXR-1 and LXRbeta in the hepatic acute phase response. *Genes Dev.* **24**, 381–395.

Wen, H., Li, J., Song, T., Lu, M., Kan, P.Y., Lee, M.G., Sha, B., and Shi, X. (2010). Recognition of histone H3K4 trimethylation by the plant homeodomain of PHF2 modulates histone demethylation. *J. Biol. Chem.* **285**, 9322–9326.

Wu, S., Wang, W., Kong, X., Congdon, L.M., Yokomori, K., Kirschner, M.W., and Rice, J.C. (2010). Dynamic regulation of the PR-Set7 histone methyltransferase is required for normal cell cycle progression. *Genes Dev.* **24**, 2531–2542.